
## **Historic, Archive Document**

Do not assume content reflects current scientific knowledge, policies, or practices.



1.0422 C4518

UNITED STATES DEPARTMENT OF AGRICULTURE PRODUCTION AND MARKETING ADMINISTRATION COTTON BRANCH BAKERSFIELD, CALIFORNIA

RECEIVED

JUL 27 1951

VOLUME XX, No. 6

DECEMBER 9, 1946

COTTON QUALITY REPORT FOR GINNINGS THROUGH NOVEMBER 30, 1946

AMERICAN-EGYPTIAN COTTON (SXP)

AMERICAN-EGYPTIAN (SXP) GINNINGS IN THE UNITED STATES TOTALED 1,936 BALES ACCORDING TO THE BUREAU OF THE CENSUS. THE DEPARTMENT OF AGRICULTURE REPORTS THE CURRENT CROP TO CONTAIN A LARGER PROPORTION OF THE HIGHER GRADES AND LONGER STAPLES THAN GINNINGS OF THIS GROWTH THROUGH THE SAME DATE IN 1945.

ARIZONA-CALIFORNIA: AMERICAN-EGYPTIAN (SXP) GINNINGS IN ARIZONA-CALIFORNIA PRIOR TO DECEMBER 1, 1946 CONTAINED A LARGER PROPORTION OF HIGHER GRADES AND LONGER STAPLES, BUT COVERED A WIDER GRADE RANGE THAN GINNINGS TO THE SAME DATE LAST YEAR. APPROXIMATELY 41 PERCENT OF THE CURRENT CROP WAS GRADES 1 AND 1-1/2, ABOUT 46 PERCENT WAS GRADES 2 AND 2-1/2 AND ALMOST 13 PERCENT WAS GRADE 3 AND LOWER. LAST SEASON 34 PERCENT WAS GRADES 1 AND 1-1/2, A LITTLE UNDER 66 PERCENT WAS GRADES 2 AND 2-1/2 AND ONLY A NEGLIGIBLE PROPORTION WAS AS LOW AS GRADE 3. STAPLES 1-1/2"AND LONGER ACCOUNTED FOR APPROXIMATELY 94 PERCENT OF THE CURRENT CROP AGAINST ABOUT 65 PERCENT A YEAR EARLIER.

NEW MEXICO-DISTRICT 6 OF TEXAS: AMERICAN-EGYPTIAN (SXP) GINNINGS IN NEW MEXICO-DISTRICT 6 OF TEXAS THROUGH NOVEMBER 30 THIS SEASON SHOWED ABOUT THE SAME PROPORTION OF THE TWO TOP GRADES AS A YEAR AGO, HOWEVER, THE CURRENT CROP CONTAINED A SUBSTANTIALLY SWALLER PROPORTION OF LONGER STAPLES. APPROXIMATELY 90 PERCENT OF THIS SEASON'S GINNINGS WAS GRADES I AND 1-1/2 AND \$BOUT IC PERCENT WAS GRADES 2 AND 2-1/2. This compares with about 92 percent and 8 percent respectively for these grade CATEGORIES LAST SEASON. ONLY ABOUT 73 PERCENT OF THE CURRENT CROP THROUGH NOVEMBER WAS 1-1/2" AND LONGER WHILE ALMOST 95 PERCENT FELL WITHIN THAT CATEGORY IN THE CORRESPONDING PERIOD LAST SEASON.

VCLUME XX NO. 6	ESTIMATED GRAI	GRADE AND STAPLE LENGTH OF AMERICAN-ESYPTIAN (SXP) COTTON GINNED	E LENGTH	OF AMERI	CAN-EGYPT	IAN (SX	OCTTON (	GINNED		DECEMBER 9, 1946	1946
	ARIZONA-CAL IFORNIA	IFORMIA		NEW MEXICO &	CO & DIST	DIST, 6 OF TEXAS	TEXAS		_T0_	TOTAL	
GRADE AND STAPLE LENGTH	NOVEMBER 14 NOVEMBER 30	SEASCF THRU NOVEMBER 30	RU 30	NO VEMBER 3	R 3 - 4	SEASON THRU NOVEMBER 30	THRU ER 30	NOVEMBER NOVEMBER	R 30	SEASON THRU	FHRU R 30
GRADE	1945 1946 PCT. PCT.	1945 PCT.	1946 PCT.	1945 PCT.	1946 PCT.	1945 PCT.	1946 PCT.	1945 PCT.	1946 PCT.	1045 PCT.	1946 <u>PCT</u> .
2-1/2 2-1/2 3-1/2	7.6 70.7 20.7 8 29.0	2012 2012 2012 2013	2007-0 2007-0 2007-0-1	448.3	20.9	37.2	37.6	8.45	23-23- 23-22-23 23-23-23-23 23-23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23-23 23-23-23-23 23-23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23-23 23-23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23-23-23 23	1,100 1,000	286- 286- 786- 787-5-
4-1/2	•		î						<b>.</b>		ν.
ALL GRADES	100.0 100.0	100.0	0.001	100.0	0.001	100.0	0.001	100.0	0.001	100.0	0.001
LENGTH (INCHES) 1-9/32 AND SHORTER. 1-5/16 AND 1-11/32. 1-7/16 AND 1-15/32. 1-1/2 AND 1-17/32. 1-9/16 AND LONGER.	: 46.1 : 53.9 94.5		2.0 2.0	23.2	6.7 47.8 45.5	5.1	4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	40.9	2.6 7.13	2.4 2.4 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0	2.2 7.4.0 4.0 4.0

\*LESS THAN 0.05 PERCENT.

ALL LENGTHS

100.0 100.0 100.0 100.0 100.0 100.0 100.0 100.0 1100.0 100.0 100.0 100.0